



京天成生物技术（北京）有限公司

AbMax Biotechnology Co., LTD

北京海淀区上地东路 29 号 2B 702

Building-2B, Suite 702, 29 E ShangDi Road,

Beijing, 100085, China

Tel: 86-10-82896224, Fax: 86-10-62977292

E-mail: info@antibodychina.com

www.antibodychina.com

Anti-Rat Cystatin C, Clone 2C2

(mouse monoclonal IgG)

Catalog # 05-0035

Lot # 100067

Immunogen: Recombinant rat Cystatin C.

Specificity: Recognizes Cystatin C. Cystatin C is known as Biomarker for kidney damage.

Species Cross-reactivity: Cross-react with human Cystatin C. Other species not determined.

Formulation: 0.2mg of mouse monoclonal IgG in 0.2ml of 0.01M Tris-HCl, pH 8.0, 0.15M NaCl, 0.02% sodium azide. Protein was determined by OD₂₈₀ absorbance.

Physical State: Frozen liquid

Storage: Store the product at -20 °C . Product is stable for about 6 weeks at 2-8 °C as an undiluted liquid. Prepare working dilution fresh each day. Avoid repeated freezing and thawing.

FOR IN VITRO RESEARCH USE ONLY

NOT FOR USE IN HUMANS

Research Applications

Indirect ELISA: A working concentration of 0.5~1µg/ml is suggested. Detection sensitivity 0.8ng/ml.

Sandwich ELISA: Used as capturing Antibody, a working concentration of 5~10µg/ml is suggested.

Immunoblot Analysis: 0.5~2µg/ml.
